


(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
14 November 2002 (14.11.2002)

PCT

(10) International Publication Number  
WO 02/090318 A1

- (51) International Patent Classification: C07C 227/12, 229/28 (74) Agents: STUART, Peter, G. et al.; Pfizer Limited, Ramsgate Road, Sandwich, Kent, CT13 9NJ (GB).
- (21) International Application Number: PCT/IB02/01401 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 18 April 2002 (18.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0110935.4 4 May 2001 (04.05.2001) GB (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): BLAKEMORE, David, Clive [GB/GB]; Pfizer Global Research and Development, Ramsgate Road, Sandwich, Kent, CT13 9NJ (GB). BRYANS, Justin, Stephen [GB/GB]; Pfizer Global Research & Development, Ramsgate Road, Sandwich, Kent, CT13 9NJ (GB).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PROCESS FOR PREPARING BICYCLIC AMINO ACIDS



(57) Abstract: The invention relates to a process for preparing (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-amino-methyl-bicyclo[3.2.0]hept-3-yl)-acetic acid, or an acid addition salt thereof. An embodiment of the process is shown in the following reaction scheme:

BEST AVAILABLE COPY

WO 02/090318 A1

# PROCESS FOR PREPARING BICYCLIC AMINO ACID


## FIELD OF THE INVENTION

5 This invention relates to a process for preparing a bicyclic amino acid, and more particularly to a process for preparing (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )-(3-aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid, or an acid addition salt thereof.

## BACKGROUND TO THE INVENTION


10

Gabapentin (Neurontin®) is an anticonvulsant agent that is useful in the treatment of epilepsy and that has recently been shown to be a potential treatment for neurogenic pain. It is 1-(aminomethyl)-cyclohexylacetic acid of structural formula:



15




Patent Application No. US 60/160725 describes a series of novel bicyclic amino acids which are analogues of gabapentin, their pharmaceutically acceptable salts, and their prodrugs of formulae:



20

wherein n is an integer of from 1 to 4. Where there are stereocenters, each center may be independently R or S, preferred compounds being those of Formulae I-IV above in which n is an integer of from 2 to 4. The compounds are disclosed as being useful in treating a variety of disorders including epilepsy, faintness attacks,

hypokinesia, cranial disorders, neurodegenerative disorders, depression, anxiety, panic, pain, neuropathological disorders, and sleep disorders. Certain of the compounds disclosed in that patent application have high activity as measured in a radioligand binding assay using [ $^3\text{H}$ ]gabapentin and the  $\alpha_2\delta$  subunit derived from porcine brain tissue (Gee N.S., Brown J.P., Dissanayake V.U.K., Offord J., Thurlow R., Woodruff G.N., *J. Biol. Chem.*, 1996;271:5879-5776). Results for some of the compounds are set out in the following table:

TABLE 1		
Compound	Structure	$\alpha_2\delta$ binding affinity ( $\mu\text{M}$ )
(1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-Aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid		0.038
(+/-)-(1 $\alpha$ ,5 $\beta$ )(3-Aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid		2.86
(1 $\alpha$ ,3 $\beta$ ,5 $\alpha$ )(3-Aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid		0.332

10

The present invention is concerned with the production of the active compound (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid, or an acid addition salt thereof. The synthetic route described in US 60/160725 proceeds via a nitro derivative produced using nitromethane, and results in a 95:5 mixture of diastereoisomers (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ ) and (1 $\alpha$ ,3 $\beta$ ,5 $\alpha$ ) respectively. The present

15


invention addresses the problem of obtaining an improved yield of product and producing a single diastereomeric product. This problem is solved by the process defined below.

5

## SUMMARY OF THE INVENTION


The present invention provides a process for preparing (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid, or an acid addition salt thereof, which comprises the following steps:

- 10 (i) condensing a cyclic ketone (1) with an alkyl cyanoacetate to form a cyanoester (2);




in which R is an alkyl group having 1 to 6 carbon atoms;

- 15 (ii) reacting the cyanoester (2) with an arylalkyl or alkenyl Grignard reagent to form a cyanoester (3):




in which R' is a phenyl or phenyl-C<sub>1</sub>-C<sub>4</sub> alkyl group or a C<sub>2</sub>-C<sub>6</sub> alkenyl group;

- (iii) removing the cyano group of the cyanoester (3) by reaction with a base to form a carboxylic acid (4):



20


- (iv) converting the carboxylic acid (4) to its alkyl ester (5):




in which R'' is an alkyl group having 1 to 6 carbon atoms;

4

(v) oxidising the alkyl ester (5) to form the acid (6):




(vi) converting the carboxylic acid group of the acid (6) to an isocyanate group, thereby forming the compound (7):




and

(vii) hydrolysing the isocyanate and ester groups of compound (7) to form the desired compound (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid (8), or an acid addition salt thereof:



### DETAILED DESCRIPTION

The starting material in the process of the invention is the cyclic ketone of formula (1). Our copending application (our reference: A0000507), the disclosure of which is hereby incorporated by reference, describes a process for preparing this cyclic ketone according to the following reaction Scheme 1:



- (I)  $\text{LiAlH}_4$ , THF, Reflux (80%);  
 (II)  $\text{MsCl}$ ,  $\text{NEt}_3$ , DCM,  $-40^\circ\text{C}$  to RT (80%);  
 (III)  $\text{NaI}$ , Acetone, Reflux (70%);  
 (IV)  $t\text{-BuLi}$ , pentane-ether (3:2),  $-25^\circ\text{C}$ ;  
 (V)  $\text{NaIO}_4$ ,  $\text{RuCl}_3 \cdot \text{H}_2\text{O}$ , MeCN, EtOAc,  $\text{H}_2\text{O}$ ;  
 (VI) MeOH, Conc  $\text{H}_2\text{SO}_4$  (85% from di-iodide);  
 (VII)  $\text{KOt-Bu}$ , THF, Reflux (97%);  
 (VIII) DMSO,  $\text{H}_2\text{O}$ ,  $155^\circ\text{C}$  (97%).

Scheme 1

Another method of preparing the cyclic ketone (1) is disclosed in  
 5 US 60/160725 and is reproduced below in Reference Example 1.

In step (i) of the process according to the invention (cf. Scheme 2 below),  
 the ketone (1) is condensed with an alkyl cyanoacetate, for example ethyl  
 cyanoacetate, preferably in an organic solvent such as toluene, benzene, xylenes  
 10 or n-heptane, to which acetic acid and  $\beta$ -alanine or ammonium acetate, or  
 piperidine are added.

Step (ii) involves the use of an arylalkyl or alkenyl Grignard reagent, and  
 results in the production of a 1:1 mixture of diastereomeric cyanoesters (3). The  
 15 arylalkyl Grignard reagent is preferably a benzyl Grignard reagent, such as  
 benzylmagnesium chloride, benzylmagnesium bromide or benzylmagnesium

iodide. Reaction with the arylalkyl Grignard reagent can be carried out at a temperature from  $-100^{\circ}\text{C}$  to  $110^{\circ}\text{C}$ , generally at room temperature.

The alkenyl Grignard reagent which may be used in step (ii) is preferably a vinyl, allyl or 2-butenyl Grignard reagent, such as vinylmagnesium chloride, vinylmagnesium bromide, allylmagnesium chloride, allylmagnesium bromide or 2-butenylmagnesium chloride. An organometallic reagent such as vinyl lithium can similarly be used. The reaction of step (ii) with an alkenyl Grignard reagent is preferably carried out in the presence of a dialkylzinc, such as dimethyl zinc, or a copper (I) salt, such as copper (I) iodide or copper (I) cyanide. This reaction is preferably carried out with cooling, for example at a temperature of from  $-100^{\circ}\text{C}$  to  $0^{\circ}\text{C}$ .

In step (iii) the cyanoester (3) is reacted with a base to remove the cyano group and hydrolyse the ester group, resulting in the single diastereomeric acid (4). The base used may be an alkali metal hydroxide, such as potassium hydroxide, sodium hydroxide, lithium hydroxide or cesium hydroxide. The reaction may be carried out in a solvent such as ethylene glycol, 2-methoxyethyl ether, 1,4-dioxane or diethylene glycol.

The carboxylic acid group of acid (4) is protected by conversion to its alkyl ester (5). The alkyl ester is preferably a methyl ester, and to obtain this the acid (53) may be added

- to a mixture of iodomethane in a solvent selected from dichloromethane, chloroform, tetrahydrofuran, toluene or 1,4-dioxane to which a base such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), triethylamine or 1,5-diazabicyclo[4.3.0]non-5-ene (DBN) is added and stirred at a temperature from  $-40^{\circ}\text{C}$  to  $110^{\circ}\text{C}$ ; or
- to a mixture of methanol and a concentrated acid such as sulphuric acid or hydrochloric acid at a temperature ranging from  $0^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ ; or

- to trimethylsilyldiazomethane and methanol in benzene or toluene at a temperature from  $-40^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ ; or
- to diazomethane in a solvent such as benzene, toluene, dichloromethane at a temperature from  $-40^{\circ}\text{C}$  to  $40^{\circ}\text{C}$ .

5        In step (v) the aryl, e.g. phenyl, group or the alkenyl, e.g. allyl, group of the resulting ester (5) is oxidized to a carboxylic acid group, for example by treatment with sodium periodate and ruthenium (III) chloride in a mixture of carbon tetrachloride or ethyl acetate and acetonitrile to which water is added. The mixture is stirred at a temperature from  $-40^{\circ}\text{C}$  to  $80^{\circ}\text{C}$  to give the carboxylic acid (6).


10

In step (vi) the carboxylic acid group of acid (6) is converted to isocyanate by addition

- to a mixture of a base selected from triethylamine or diisopropylethylamine and a solvent selected from toluene, benzene, 15 xylenes, tetrahydrofuran, diethyl ether or *n*-heptane to which diphenylphosphoryl azide (DPPA) is added and stirring at a temperature from  $0^{\circ}\text{C}$  to  $150^{\circ}\text{C}$  to produce the isocyanate of formula (7); or
- to ethyl chloroformate or isobutyl chloroformate and a base such as triethylamine or diisopropylethylamine in tetrahydrofuran or acetone or 20 diethyl ether at a temperature of  $-40^{\circ}\text{C}$  to  $78^{\circ}\text{C}$  followed by addition of sodium azide in water and tetrahydrofuran or acetone followed by addition of toluene or benzene and refluxing.

25        In step (vii), the isocyanate and ester groups of compound (7) are simultaneously hydrolysed to amino and carboxylic acid groups, e.g. by aqueous hydrochloric acid at a concentration of from 0.01 M to 12 M optionally in the presence of a solvent such as 1,4-dioxane, acetic acid or water to produce the amino acid (8).

An embodiment of the invention using a benzyl Grignard reagent in step (ii) is detailed below ("benzyl route"). The main advantage of this route is that the addition of the benzyl Grignard reagent  $\text{BnMgCl}$  can be carried out at room temperature without the necessity for an additive (such as dimethylzinc or copper (I) cyanide). The benzyl Grignard addition also appears to be stereoselective (there being no evidence from NMR or GC analysis for the presence of more than one diastereoisomer of the benzyl acid after hydrolysis of the cyanoester).




- (i)  $\text{NCCH}_2\text{CO}_2\text{Et}$ ,  $\text{NH}_4\text{OAc}$ ,  $\text{AcOH}$ , Toluene, Reflux;
- (ii)  $\text{BnMgCl}$ , THF,  $15^\circ\text{C}$ ;
- (iii)  $\text{KOH}$ , Ethylene glycol,  $160^\circ\text{C}$  (95% from Knoevenagel product);
- (iv)  $\text{TMSCHN}_2$ , Toluene (92% from cyanoester);
- (v)  $\text{RuCl}_3$ ,  $\text{NaIO}_4$ ,  $\text{CCl}_4$ ,  $\text{MeCN}$ ,  $\text{H}_2\text{O}$  (66%);
- (vi)  $\text{DPPA}$ ,  $\text{NEt}_3$ , Toluene, Reflux;
- (vii)  $6\text{N HCl}$ , reflux (72% from acid).
- [ $\text{TMSCHN}_2$  = trimethylsilyldiazomethane;  $\text{DPPA}$  = diphenylphosphoryl azide]

### Scheme 2

An embodiment of part of the process of the invention in which an allyl Grignard reagent is used in step (ii) is detailed in Schemes 3 and 4 below ("allyl

route"). The main advantage of this route is that the allyl oxidation (to the carboxylic acid) requires only four equivalents of sodium periodate in addition to the ruthenium trichloride. The main disadvantage of this route is that the conjugate addition of the allyl Grignard requires an additive such as dimethylzinc or copper (I) cyanide. The yields obtained with dimethylzinc over the two steps of conjugate addition and hydrolysis were higher than with the cuprate addition (89% as opposed to 70%).




(ii) Allylmagnesium bromide, dimethylzinc, THF,  $-78^{\circ}\text{C}$ ;  
(iii) KOH, Ethylene glycol,  $165^{\circ}\text{C}$  (89% over the two steps)

OR

(ii) Allylmagnesium bromide,  $\text{Cu(I)CN}$ , THF,  $-78^{\circ}\text{C}$ ;  
(iii) KOH, Ethylene glycol,  $165^{\circ}\text{C}$  (69% over the two steps)

Scheme 3

The copper (I) cyanide reaction was investigated at two different temperatures and while the reaction appears to go cleanly at 0 °C (apparently giving a single diastereoisomer after hydrolysis) the yield was poorer than at -78 °C (presumably due to polymerisation). However, other temperatures may be used.



(iv) TMSCHN<sub>2</sub>, Toluene (97% from cyanoester);

(v) RuCl<sub>3</sub>, NaIO<sub>4</sub>, CCl<sub>4</sub>, MeCN, H<sub>2</sub>O (83%);

(vi) DPPA, NEt<sub>3</sub>, Toluene, Reflux;

(vii) 6N HCl, reflux (72% from acid).


#### Scheme 4

The invention is illustrated by the following Examples.

#### EXAMPLES

15

##### REFERENCE EXAMPLE 1



**Synthesis of compound A**

Lithium aluminum hydride (69.4 mL of a 1 M solution in ether, 69.4 mmol) was added dropwise to a stirring solution of *cis*-cyclobutane-1,2-dicarboxylic acid (5 g, 34.7 mmol) in THF (60 mL) at 0°C under argon. The mixture was allowed to warm to room temperature and stirred for 16 hours. The mixture was cooled to 0°C and quenched by careful addition of water (2.7 mL), sodium hydroxide solution (2.7 mL of a 15% w/v solution), and water (8.1 mL). The mixture was stirred for 15 minutes, and the precipitate was removed by filtration. The solvent was evaporated under reduced pressure to give the *alcohol* A as a colorless oil (4.0 g, 98%);  $\delta_{\text{H}}$  (400 MHz;  $\text{CDCl}_3$ ): 3.85 (2H, m), 3.6 (2H, m), 3.2 (2H, s), 2.7 (2H, m), 2 (2H, m); 1.55 (2H, m);  $\delta_{\text{C}}$  (400 MHz;  $\text{CDCl}_3$ ): 63.15, 37.83, 20.40.

**Synthesis of compound B**

Mesyl chloride (6.2 mL, 79.1 mmol) was added dropwise to a stirring solution of A (4.0 g, 34.4 mmol) in dichloromethane (150 mL) at -40°C under argon. Triethylamine (12.0 mL, 86.0 mmol) was then added dropwise, and the mixture was allowed to warm slowly to room temperature. After stirring for 16 hours, the mixture was quenched by addition of dilute hydrochloric acid (50 mL). The organic layer was separated, and the aqueous layer was further extracted with dichloromethane (2  $\times$  50 mL). The combined organic fractions were washed with brine, dried ( $\text{MgSO}_4$ ), and the solvent was evaporated under reduced pressure. The residue was chromatographed ( $\text{SiO}_2$ , heptane/ethyl acetate, 6:4) to give the *mesylate* B (6.1 g, 73%) as a white solid;  $R_{\text{f}}$  (heptane/ethyl acetate, 1:1) 0.18.  $\delta_{\text{H}}$  (400 MHz;  $\text{CDCl}_3$ ): 4.3 (4H, m), 3.05 (6H, s), 2.9 (2H, m), 2.2 (2H, m), 1.8 (2H, m);  $\delta_{\text{C}}$  (400 MHz;  $\text{CDCl}_3$ ): 69.51, 37.45, 35.28, 21.09.

**Synthesis of compound C**

Anhydrous lithium bromide (10.6 g, 121.8 mmol) was added to a stirring mixture of B (5.95 g, 24.4 mmol) in acetone (50 mL) under argon and the mixture was refluxed for 2 hours. After cooling, the acetone was evaporated under reduced pressure and the residue was taken up in ether (50 mL), washed with water


(50 mL), brine, dried ( $\text{MgSO}_4$ ), and the solvent was evaporated under reduced pressure. The residue was chromatographed ( $\text{SiO}_2$ , heptane/ethyl acetate, 95:5) to give the *dibromide* C (5.36 g, 86%) as an orange liquid;  $R_f$  (heptane-ethyl acetate, 8:2), 0.82.  $\delta_H$  (400 MHz;  $\text{CDCl}_3$ ): 3.6 (2H, m), 3.45 (2H, m), 2.85 (2H, m), 2.1 (2H, m), 1.7 (2H, m;  $\delta_C$  (400 MHz;  $\text{CDCl}_3$ ): 39.70, 33.79, 23.95.

#### Synthesis of compound 1

To a cooled ( $0^\circ\text{C}$ ) suspension of potassium hydride (1.58 g, 39.5 mmol) (previously washed 3 times with pentane) in tetrahydrofuran (22 mL) was added, under an argon atmosphere, a solution of methyl methylthiomethyl sulfoxide (1.36 mL, 13.04 mmol, previously dried over molecular sieves for 3 hours) in tetrahydrofuran (3 mL) over 1 hour. After stirring for a further 30 minutes, a solution of C (3.17 g, 13.1 mmol) in THF (2 mL) was added, at  $0^\circ\text{C}$ , over 1 hour. The reaction mixture was then allowed to warm up to room temperature and was stirred overnight. The mixture was quenched by addition of aqueous ammonium chloride (6 mL, 25%). After 10 minutes, the solid was filtered off and the filtrate concentrated. The residue was taken up in ether (20 mL) and 9N sulfuric acid (0.05 mL) was added. After stirring for 30 hours, saturated sodium hydrogen carbonate was added. The ether phase was separated and concentrated to 5 mL. Saturated sodium hydrogen sulphite (1.5 g) solution was added and the mixture stirred for 30 minutes. The phases were separated. The ethereal phase was stirred for further 30 minutes with a saturated sodium hydrogen sulphite (0.5 g) solution. The phases were separated and the collected aqueous phases were treated with aqueous sodium hydroxide (5 mL, 20%) and extracted with ether. The ether phase was dried ( $\text{MgSO}_4$ ) and evaporated under reduced pressure to give 4 as a yellow liquid (0.16 g, 11%).  $\delta_H$  (400 MHz;  $\text{CDCl}_3$ ): 3.0 (2H, m), 2.15-2.45 (6H, m), 1.65 (2H, m).

## EXAMPLE 1

Synthesis of:




5 Ketone (1) (199.3 mmol), ethyl cyanoacetate (21.2 ml, 199.3 mmol), ammonium acetate (15.4 g, 199.3 mmol) and glacial acetic acid (11.4 ml, 199.3 mmol) were refluxed in toluene (250 ml) using a Dean-Stark trap. After 8 h, the mixture was allowed to cool and diluted with ethyl acetate (400 ml), washed with water (3 x 150 ml), brine and dried (MgSO<sub>4</sub>). The solvent was evaporated under  
10 reduced pressure. The residue was chromatographed (SiO<sub>2</sub>, heptane/ethyl acetate, 95:5 to 7:3) to give *cyano-ester* (2) (31.95g, 78%); R<sub>f</sub> (heptane-ethyl acetate, 8:2) 0.40;  $\nu_{\max}(\text{film})/\text{cm}^{-1}$  2226 (CN), 1727 (C=O), 1614 (C=C);  $\delta_{\text{H}}(400 \text{ MHz}; \text{CDCl}_3)$  4.29 (2H, q, *J* 7.1, CO<sub>2</sub>CH<sub>2</sub>Me), 3.34 (1H, d, *J* 20), 3.08-2.88 (5H, m), 2.30-2.18 (2H, m), 1.60-1.51 (2H, m), 1.36 (3H, t, *J* 7.1, CO<sub>2</sub>CH<sub>2</sub>Me); *m/z* (CI) 204 (M-H,  
15 100%).

## EXAMPLE 2

Synthesis of:

20




Cyanoester (2) (12.0 g, 59 mmol) in THF (50 ml) was added over 2 h to a stirring solution of benzylmagnesium chloride (117 ml of a 1M solution in ether, 117 mmol) in THF (300 ml) at 15°C under argon. After allowing the solution to  
25 warm to room temperature the mixture was stirred for a further 1 h and then the

mixture was cooled to 15°C and quenched by addition of saturated ammonium chloride solution (100 ml). The mixture was allowed to warm to room temperature, and dilute hydrochloric acid (300 ml) was added. The aqueous layer was extracted with ethyl acetate (2 x 300 ml). The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>) and the solvent was evaporated under reduced pressure to give a 1:1 mixture of diastereomeric *cyano-esters* (3) (19.85g, > 100 % crude yield); R<sub>f</sub>(heptane-ethyl acetate, 9:1) 0.25;  $\nu_{\max}(\text{film})/\text{cm}^{-1}$  2246 (CN), 1741 (C=O); *m/z* (CI) 296 (M-H, 100%); (CI<sup>+</sup>) 298 (M+H, 90%).

10

**EXAMPLE 3**

Synthesis of:




The mixture of diastereomeric *cyano-esters* (3) (17.39 g, 59 mmol) and potassium hydroxide (19.67 g, 351 mmol) were heated to 160 °C in ethylene glycol (400 ml) for 38 h. After this time, the mixture was allowed to cool and dilute hydrochloric acid (300 ml) was added carefully. The mixture was extracted with ethyl acetate (3 x 200 ml) and the combined organic fractions were washed with brine, dried (MgSO<sub>4</sub>) and the solvent was evaporated under reduced pressure. The residue was chromatographed (SiO<sub>2</sub>, heptane/ethyl acetate, 8:2) to give the single diastereomeric *acid* (4) (15.96 g, 95 %); R<sub>f</sub>(heptane-ethyl acetate, 1:1) 0.67;  $\nu_{\max}(\text{film})/\text{cm}^{-1}$  1703 (C=O);  $\delta_{\text{H}}(400 \text{ MHz}; \text{CDCl}_3)$  7.30-7.17 (5H, m, Ph), 2.84 (2H, m), 2.55 (2H, s, CH<sub>2</sub>Ph), 2.44 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.29 (2H, m), 2.02 (2H, dd, *J* 13.2, 8.3), 1.66 (2H, m), 1.50 (2H, dd, *J* 12.7, 5.2); *m/z* (CI) 243 (M-H, 55%);

15

## EXAMPLE 4

Synthesis of:

5




Trimethylsilyldiazomethane (43.2 ml of a 2M solution in hexane, 86.4 mmol) was added dropwise to a stirring solution of acid (4) (17.55 g, 72 mmol) in a mixture of toluene (120 ml) and methanol (50 ml) at 10 °C under argon over 30 minutes. The mixture was allowed to warm to room temperature and stirred for 1 h. The solvent was removed under reduced pressure and the residue was taken up in ethyl acetate (300 ml), washed with saturated sodium hydrogen carbonate (300 ml), dilute hydrochloric acid (300 ml), brine and dried (MgSO<sub>4</sub>). The solvent was evaporated under reduced pressure to give *ester* (5) (17.08 g, 92%); *R*<sub>f</sub>(heptane-ethyl acetate, 9:1) 0.51;  $\nu_{\max}(\text{film})/\text{cm}^{-1}$  1737 (C=O);  $\delta_{\text{H}}(400 \text{ MHz}; \text{CDCl}_3)$  7.39-7.15 (5H, m, Ph), 3.71 (3H, s, OMe), 2.81 (2H, m), 2.51 (2H, s, CH<sub>2</sub>CO<sub>2</sub>Me), 2.39 (2H, s, CH<sub>2</sub>Ph), 2.26 (2H, m), 1.97 (2H, dd, *J* 13.3, 8.4), 1.64 (2H, m), 1.47(2H, dd, *J* 12.5, 5.1).

20

## EXAMPLE 5

Synthesis of:




25

Ester (5) (10.08 g, 39 mmol) and sodium periodate (117 g, 55 mmol) were stirred together in ethyl acetate (58 ml), acetonitrile (58 ml) and water (512 ml)

for 5 minutes. The mixture was cooled to 0 °C and ruthenium (III) chloride (0.162 g, 0.8 mmol) was added to the reaction mixture. The reaction was allowed to warm to room temperature and stirred for 24 h with intermittent cooling with an ice bath to control the exotherm. Diethyl ether (100 ml) was added and the mixture was stirred for 40 minutes. The mixture was poured onto dilute hydrochloric acid and extracted with ethyl acetate (2 x 300 ml). The combined organic fractions were washed with brine, dried (MgSO<sub>4</sub>) and the solvent was evaporated under reduced pressure. The residue was purified by chromatography (SiO<sub>2</sub>, heptane to 8:2 heptane/ethyl acetate) to give the *acid* (6) (6.21 g, 66.2 %);  
10  $R_f$ (heptane-ethyl acetate, 1:1) 0.47;  $\nu_{\max}$ (film)/cm<sup>-1</sup> 1737 (C=O), 1704 (C=O);  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 3.71 (3H, s, OMe), 2.80-2.71 (4H, m), 2.33 (2H, s), 2.26 (2H, m), 2.07 (2H, m), 2.05 (2H, s), 1.64 (2H, m), 1.54 (2H, dd,  $J$  13.2, 5.2);  $m/z$  (CI) 225 (M-H), (CIP<sup>+</sup>) 227 (M+H). -

15


**EXAMPLE 6****Synthesis of:**

Diphenylphosphoryl azide (3.66 g, 17 mmol), triethylamine (2.43 g, 17.5 mmol), and acid (6) (3.8 g, 16.8 mmol) were refluxed in toluene (50 ml) for 1.25 h. The mixture was allowed to cool and diluted with ethyl acetate (200 ml). The resulting solution was washed with saturated aqueous sodium hydrogen carbonate (200 ml), brine, and dried (MgSO<sub>4</sub>). The solvent was removed under reduced pressure to give the *isocyanate* (7) which was used without further purification  
25 (3.75 g, 100%);  $R_f$ (heptane-ethyl acetate, 9:1) 0.39;  $\nu_{\max}$  (film)/cm<sup>-1</sup> 2266 (NCO), 1733 (C=O);  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 3.69 (3H, s, OMe), 3.17 (2H, s, CH<sub>2</sub>NCO), 2.69 (2H, m), 2.58 (2H, s, CH<sub>2</sub>CO<sub>2</sub>Me), 2.24 (2H, m), 1.94 (2H, m), 1.65 (2H, m), 1.56 (2H, dd,  $J$  12.9, 4.6).

## EXAMPLE 7

Synthesis of:

5




The isocyanate (7) (9.88 g, 45 mmol) and 6N hydrochloric acid (100 ml) were refluxed at 130 °C for 18 h. The mixture was allowed to cool, diluted with water (200 ml) and extracted with dichloromethane (2 x 200 ml). The aqueous  
10 was concentrated to an orange solid and washed with acetonitrile to give the *hydrochloride salt* of compound (8) (7.10 g, 73%);  $\delta_H$ (400 MHz;  $d_6$ -DMSO) 7.88 (2H, br s, NH<sub>2</sub>), 2.67 (4H, s), 2.60 (2H, s), 2.22-2.11 (2H, m), 1.94 (2H, dd, *J* 13.5, 8.0), 1.62 (2H, m), 1.52 (2H, dd, *J* 13.5, 4.9); *m/z* (ES<sup>+</sup>) 184 (M+H, 100%); LCMS (Prodigy ODS3 (3 $\mu$ ) 150 mm x 4.6 mmid column, 20-100% Acetonitrile +  
15 0.1% formic acid) Retention Time = 4.44 min, 100% purity.

The following Examples 8, 9 and 10 illustrate the use of an allyl Grignard reagent in step (ii) and thus involve an allyl compound in steps (iii) and (iv) ("allyl route").

20

## EXAMPLE 8

Synthesis of:



25

Compound (3A) can be made in two different ways and this affects the yield of compound (4A) as purification is not carried out after synthesis of (3A):

*Method A (copper (I) cyanide route)*

Allylmagnesium bromide (32.2 ml of a 1 M solution in diethyl ether, 32.2 mmol) was added dropwise to a stirring suspension of copper (I) cyanide (1.44 g pre-dried under vacuum, 16.1 mmol) in THF (60 ml) at 0 °C under argon. After 45 mins, the mixture was cooled to -78 °C and cyanoester (2) (produced as in Example 1) (3.0 g, 14.62 mmol) in THF (30 ml) was added over 1 h. After stirring for a further 1 h, the mixture was quenched by addition of saturated basic ammonium chloride (50 ml of a solution of saturated ammonium chloride with 10% [v/v] concentrated ammonia added). After warming to room temperature, diethyl ether (200 ml) was added and the organic layer was further washed with saturated basic ammonium chloride (3 x 100 ml) until the aqueous layer was no longer blue. The organic layer was washed with brine, dried (MgSO<sub>4</sub>) and the solvent was removed under reduced pressure to give a mixture of diastereomeric mixture of *cyanoesters* (3A);  $R_f$ (heptane-ethyl acetate, 7:3) 0.54;  $\nu_{\max}(\text{film})/\text{cm}^{-1}$  2247 (CN), 1742 (C=O);  $m/z$  (CI) 246 (M-H, 100%).

The mixture of diastereomeric cyano-esters (3A) (approx 14.6 mmol) and potassium hydroxide (4.92 g, 87.7 mmol) were heated to 160 °C in ethylene glycol (400 ml) for 4 days. After this time, the mixture was allowed to cool and dilute hydrochloric acid (300 ml) was added carefully. The mixture was extracted with ethyl acetate (3 x 200 ml) and the combined organic fractions were washed with brine, dried (MgSO<sub>4</sub>) and the solvent was evaporated under reduced pressure. The residue was chromatographed (SiO<sub>2</sub>, heptane/ethyl acetate, 8:2) to give the single diastereomeric *acid* (4A) (1.97g, 69%);  $R_f$ (heptane-ethyl acetate, 1:1) 0.67;  $\nu_{\max}(\text{film})/\text{cm}^{-1}$  1705 (C=O);  $\delta_H$ (400 MHz; CDCl<sub>3</sub>) 5.78 (1H, ddt,  $J$  17.1, 10.0, 7.6, CH<sub>2</sub>CH=CH<sub>A</sub>H<sub>B</sub>), 5.09-4.98 (2H, m, CH<sub>2</sub>CH=CH<sub>A</sub>H<sub>B</sub>), 2.69 (2H, m), 2.50

(2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 2.17 (2H, m), 2.01-1.93 (4H, m), 1.64 (2H, m), 1.53 (2H, dd,  $J$  12.8, 5.1);  $m/z$  ( $\text{C}^+$ ) 195 ( $\text{M}+\text{H}$ , 100%);

#### 5 Method B (Dimethylzinc method)


Allylmagnesium bromide (13.4 ml of a 1 M solution in diethyl ether, 13.4 mmol) was added to a stirring solution of dimethylzinc (6.7 ml of a 2M solution in toluene, 13.4 mmol) in THF (50 ml) at 0 °C under argon. After 20 mins, the mixture was cooled to -78 °C and cyanoester (2) (produced as in Example 1) (2.5 g, 12.18 mmol) in THF (30 ml) was added dropwise over 1 h. After stirring for a further 1 h, the mixture was quenched by careful addition of saturated ammonium chloride solution (30 ml). After warming to room temperature, dilute hydrochloric acid (100 ml to solubilise the zinc salts) was added and the mixture was extracted with diethyl ether (3 x 150 ml). The combined organic fractions were washed with brine, dried ( $\text{MgSO}_4$ ) and the solvent removed under reduced pressure to give the diastereomeric mixture of *cyanoesters* (3A).

The mixture of diastereomeric cyano-esters (3A) (approx 12.2 mmol) and potassium hydroxide (4.1 g, 73.1 mmol) were heated to 160 °C in ethylene glycol (400 ml) for 4 days. After this time, the mixture was allowed to cool and dilute hydrochloric acid (300 ml) was added carefully. The mixture was extracted with ethyl acetate (3 x 200 ml) and the combined organic fractions were washed with brine, dried ( $\text{MgSO}_4$ ) and the solvent was evaporated under reduced pressure. The residue was chromatographed ( $\text{SiO}_2$ , heptane/ethyl acetate, 8:2) to give the single diastereomeric *acid* (4A) (2.1g, 89%).

### EXAMPLE 9

Synthesis of:

20




Trimethylsilyldiazomethane (13 ml of a 2M solution in hexane, 25 mmol) was added dropwise to a stirring solution of acid (4A) (4.07 g, 21 mmol) in a mixture of toluene (40 ml) and methanol (10 ml) at 5 to 15°C under argon over 30 minutes. The mixture was allowed to warm to room temperature and stirred for 1 h. The solvent was removed under reduced pressure and the residue was taken up in ethyl acetate (300 ml), washed with saturated sodium hydrogen carbonate (300 ml), dilute hydrochloric acid (300 ml), brine and dried (MgSO<sub>4</sub>). The solvent was evaporated under reduced pressure to give *ester* (5A) (4.22 g, 96.5 %);

10  $R_f$ (heptane-ethyl acetate, 9:1) 0.44;  $\nu_{\max}$ (film)/cm<sup>-1</sup> 1738 (C=O);  $\delta_H$ (400 MHz; CDCl<sub>3</sub>) 5.78 (1H, ddt,  $J$  17.1, 10.0, 7.3, CH<sub>2</sub>CH=CH<sub>A</sub>H<sub>B</sub>), 5.07-4.97 (2H, m, CH<sub>2</sub>CH=CH<sub>A</sub>H<sub>B</sub>), 3.67 (3H, s, OMe), 2.69 (2H, m), 2.46 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.24 (2H, m), 1.95-1.90 (4H, m), 1.65 (2H, m), 1.50 (2H, dd,  $J$  12.5, 5.1);  $m/z$  (Cl<sup>+</sup>) 209 (M+H, 100%);

15

### Example 10

#### Synthesis of:



20 Ester (5A) (4.22 g, 20 mmol) and sodium periodate (18.23 g, 85 mmol) were stirred together in ethyl acetate (31 ml), acetonitrile (31 ml) and water (270 ml) for 5 minutes. The mixture was cooled to 5 °C and ruthenium (III) chloride (0.044 g, 0.4 mmol) was added to the reaction mixture. The reaction was allowed to warm to room temperature and stirred for 24 h with intermittent cooling with an

25 ice bath to control the exotherm. Diethyl ether (100 ml) was added and the mixture was stirred for 40 minutes. The mixture was poured onto dilute

hydrochloric acid and extracted with ethyl acetate ( 2x400 ml). The combined organic fractions were washed with brine, dried ( $\text{MgSO}_4$ ) and the solvent was evaporated under reduced pressure. The residue was purified by chromatography ( $\text{SiO}_2$ , heptane to 8:2 heptane/ethyl acetate) to give the *acid* (6) (3.8 g, 83 %).

5


The acid (6) can then be converted to the isocyanate (7) and the desired produce (8) as in Examples 6 and 7.

## CLAIMS

1 A process for preparing (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid, or an acid addition salt thereof, which comprises the following


5 steps:

(i) condensing a cyclic ketone (1) with an alkyl cyanoacetate to form a cyanoester (2):




in which R is an alkyl group having 1 to 6 carbon atoms;

10 (ii) reacting the cyanoester (2) with an arylalkyl or alkenyl Grignard reagent to form a cyanoester (3):




in which R' is a phenyl or phenyl-C<sub>1</sub>-C<sub>4</sub> alkyl group or a C<sub>2</sub>-C<sub>6</sub> alkenyl group;

15 (iii) removing the cyano group of the cyanoester (3) by reaction with a base to form a carboxylic acid (4):




(iv) converting the carboxylic acid (4) to its alkyl ester (5):




in which R'' is an alkyl group having 1 to 6 carbon atoms;

20 (v) oxidising the alkyl ester (5) to form the acid (6):




(vi) converting the carboxylic acid group of the acid (6) to an isocyanate group, thereby forming the compound (7):



and

(vii) hydrolysing the isocyanate and ester groups of compound (7) to form the desired compound (1 $\alpha$ ,3 $\alpha$ ,5 $\alpha$ )(3-aminomethyl-bicyclo[3.2.0]hept-3-yl)-acetic acid (8), or an acid addition salt thereof:



2 A process according to Claim 1, in which the arylalkyl Grignard reagent used in step (ii) is a benzyl Grignard reagent.

3 A process according to Claim 2, in which the benzyl Grignard reagent is benzylmagnesium chloride, benzylmagnesium bromide or benzylmagnesium iodide.

4 A process according to Claim 1, in which the alkenyl Grignard reagent used in step (ii) is a vinyl, allyl or 2-butenyl Grignard reagent.

5 A process according to Claim 4, in which the Grignard reagent is vinyl lithium, allylmagnesium chloride, allylmagnesium bromide or 2-butenylmagnesium chloride.

6 A process according to Claim 4 or 5, in which step (ii) is carried out in the presence of a dialkylzinc or a copper (I) salt.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 02/01401

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07C227/12 C07C229/28

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 28978 A (RECEVEUR JEAN MARIE ;BLAKEMORE DAVID CLIVE (GB); BRYANS JUSTIN STE) 26 April 2001 (2001-04-26) cited in the application page 45	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*A\* document member of the same patent family

Date of the actual completion of the international search

1 July 2002

Date of mailing of the international search report

10/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Sánchez García, J.M.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 02/01401

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0128978	A	26-04-2001 AU	1092001 A
		WO	0128978 A1
			30-04-2001
			26-04-2001

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**